

## SUPPLEMENTARY MATERIAL

**Supplementary Table 1.** ICU patient characteristics.

	Initial adequate treatment	Initial inadequate treatment
<b>Sex, n</b>		
<b>Male</b>	7	5
<b>Female</b>	2	2
<b>Age, mean</b>	56.6	63.9
<b>Treatment adequateness, n (%)</b>	9	7
<b>CRGN Pathogen, n</b>		
<i>Klebsiella pneumoniae</i>	6	2
<i>Stenotrophomonas maltophilia</i>	1	0
<i>Pseudomonas aeruginosa</i>	2	5
<b>Severity at admission qSOFA), n</b>		
qSOFA 0	4	3
qSOFA 1	4	1
qSOFA 2	0	2
qSOFA 3	1	1
<b>Suspected origin of bacteraemia, n</b>		
Pneumonia	6	1
Catheter	3	3
Intraabdominal	0	1
Other	0	2
<b>Shock during bacteraemia, n (%)</b>	7/8 (88%)*	4/7 (57%)
<b>Mechanical ventilation during bacteraemia, n (%)</b>	8/8 (100%)*	6/7 (86%)
<b>Need of dialysis due to acute renal failure, n (%)</b>	3/8 (38%)*	2/7 (29%)

\*1 patient had missing data on this items, reason why only 8 patients were included in the analysis despite being 9 patients in the group.